

```
5007时中的短期都有是一种时间的一种的是一种的一种。
                                                          3 /2816 64/000/033/0029/0033
          TENER V. I. Belenko, R. M.: Krylov, A. J.: Panferov, I. M.: Romanova,
         Marsa, I. Te.; Shiikina, 2. ..
        TResults of Satellite Observations
mmggme AN SCSP. Astronomicheskiy sovet. Byulleten' stantsiy opticheskogo marayadeniya skusstvennykh sputnikov Zemli, no. 33, 1963. 29-33
THE TABLE artificial matellite, satellite tracking, matellite tracking camera/
Bath lite 1966 ., NAPA 58/25 camera, KIM 5 microscope, Ural computer
        thervations were made on the satellite 1960 to during August and October
         St. 4 % (%) camera was used. The observer was A. J. Kry.ov. Measurements
          Wikit microscope by R. M. Belenke. L. M. Fanish v. and v. f.
           completations were made by the Klaeley method in two sats of three
          sing and by the Turner method. The Period as the transport of the anti-sins on
           opposer. In a serve tion times were reduced to standard time. The last co.-
           The element possible maximal error in occitionates to also of intellabil-
 ating optical center within 'cm. V. I. Belanks and J. J. Shikina
part, pated in the work. Results of 125 observations are given in a table, part
Card 1/5
```









Table 1 (1971) - A75(-O554)


- Size if wireliability of determining optics; benter within 1 mm. 7. 1. Selence is no payon ipated in the word, one, is in the more are given and given in a reproduce, in the limit with the latter.

- Size in the first transfer of the control of the more and the control of the more and payon. AN SSSE)


- Size in the control of the con



	·
* * * * * A *	PROFESSION AND STREET CHART
$\frac{\Lambda_{i}}{2} \frac{\Lambda_{i}}{\Lambda_{i}} \frac{\Lambda_{i}}{2} = 0$ whenever, $G_{i}$ , $G_{i}$ , $G_{i}$ , $G_{i}$	In the Continue 2, 0.199
terration of artificial satellites	• •
1 1 AN 11W Astronomicheskiy sovet. Byulleten	' stants y onto cheskom
om ynterlys lek isstvennych sputnikov čemli, no. 52	
	, 1965, CE 17
TOPIC TAGS: ertificial satellite, satellite tracki	ng, satellite tracking camera/
TOPIC TAGS: artificial satellite, satellite trackis satellite 14 11, NAFA-3s/25 camera, KIM 3 microsofth Topic Topic 14 11, NAFA-3s/25 camera, KIM 3 microsofth Topic Topic Napala NAFA-3s/25. Observers were A. G. Krylov ments were made on a KIM-3 microscope by R. E. Bele to a nature by the Kinelay mathod for two sets	ng, satellite tracking camera/ ope, Ural computer  1950 . in Tune 1911. The camera and Y. A. Yurevich. Measure- ako and I. M. Panferov. Computa-
TOPIC TAGS: artificial matellite, matellite tracking the lite of t	ng, satellite tracking camera/ ope, Ural computer  1950 . in Tune 1911. The camera and Y. A. Yurevich. Measure- ako and I. M. Panferov. Computa-
TOPIC TAGS: artificial satellite, satellite trackis satellite in, NAFA-38/25 camera, KIM 3 microsofth Class of the American were made on the satellite tracking was a NAFA-38/25. Observers were A. G. Krylov ments were made on a KIM-3 microscope by R. M. Bele to a contract of the satellite of	ng, satellite tracking camera/ ope, Ural computer  1950 in the 1961. The camera and Y. A. Yurevich. Measure- ako and I. M. Panferov. Computa- of three reference stars and by
TOPIC TAGS: ertificial matellite, matellite tracki matellite 14 11, NAFA-38/25 camera, KIM 3 microsofth Tag Topic honorvations were made on the matellite tracki was a NAFA-38/25. Observers were A. G. Krylov ments were made on a KIM-3 microscope by R. E. Bele track by the Kinelay mathod for two metals.	ng, satellite tracking camera/ ope, Ural computer  1950 . in tune 1911. The camera and Y. A. Yurevich. Measure- ako and I. M. Panferov. Computa- of three reference stars and by




	no teoreta At	<b>150</b> 0 1547			7. N.	5URB: 01	
		Station of the Astronomical Journal AN ALLE					
	н б н б	0.F.	ΔΙ	OI 4			
		<u>.</u> . L.		•			
		•					
	rma i	20h5g和56 <sup>8</sup> ,404	10 <sup>8</sup> .00€	Oth Cartain	· 0 · 8 · 83 #	9" I	
•	, i	22 29 52.794	0.00%	D2 5x 4h .4		··· II	
j							



A country are Alfoutfide

ASP CATT No stantsiya Astronomicheskogo soveta AN SSSR (Station of the Astronomical Council, AN SSSR)

UTRNITTED \*\*Dec62 ENCL: 01 SUB CODE: SV, DC NC REF SOV. COO OTHER: OCC



SEEDIN MR: ATSOC5775 \$/2616/65/000/036/0031**/0033** AUTHORS: Belenko, R. M.; Krylov, A. G.; Panferov, I. M.; Sentsova, Yu. Ye.; Shilkina, Z. S.; Yurevich, V. A. Results of Satellite Observations/ 31 日野、Astronomicheskly sovet. Byulleten' stante'v opticheskogo nablyudeniya lakusetvennyen eputnikoy Zemil, no. 35, 1963, /-53 TOPIC TAGS: artificial satellite, satellite tracking camera, satellite track analysia/ satellite 1961 & E., satellite 1961 &, satellite 1962 ( , satellite such that the tip, autolite 1960  $\xi$  . NAFA is is named at XIM  $^3$  microscope, and rose ope, Wral 1 computer Tranvations were made in April, May, and home 'm' on the satellites عدد معروب المراج المراج المراج المراج المراج المراجع ا A. G. Krylov (indicated by II in the table) and V. A. Yurevich (I in table). Both 11. 15. 16. 16. 15. Cameras. Measurements were made by R. M. Belenko with a KIM-5 . M. Panterov, with a TIM- Coil of the life of easing was line by sa toto by You Ye. Sentabya Hainkia Toki omenit in computer. Por


1.8 % - a.15003773		
Such and Military the same of	oordinates, the coordinates of one coordinates of points on the Rall table shows deviation of the coordinates given in the Rall and members reduced to the coordinates given to the Rall and the Coordinates given in the Rall and the Rall and the Rall and the Coordinates and the Coordinates and the Coordinates and the Coordinates are coordinates. No. 1072	ite oute track. The next spot the spot the spot the spot the spot the spot track in a spot the spot track in the spot track.
SUPERIOR SERVICE	ENGL: 01	JUH JODE: SY, DC
8 (EF SOV) - 200	CTHER: 000	
erw prz		
では、 では、 では、 では、 では、 では、 では、 では、		

to the American to				सम्बद्धाः । १५ सम्बद्धाः । १५
	The Astronomical C	ouncil of the A	UN SESP	
2, 1 5.		f	• •	amera
<u>1962</u>	196	19 6,		
	25 <sup>h</sup> 44 <sup>m</sup> 25 <sup>8</sup> 277 ±0.95	18 <sup>h</sup> 40 <sup>m</sup> 58 <sup>s</sup> 8	140	II
me 1.	20 44 33.193 20.01	22 21 16.1	34 51 00	II

KRYLOV, A.G.

Classification of ceder forests in the Altai. Izv. SO AN SSSR no.8 Ser. biol.-med. nauk no.2:40-47 64 (MIRA 18:1)

1. Institut lesa i drevesiny Sibirskogo otdeleniya AN SSSR, Krasnoyarsk.



	.R: AT5012011			İ
i kaleng d	api tee Kurile e e u	s, and maintenance of well Islands. The original color of the color of	i a venere e	
	income .			
and Allenia	A: Astronomiches	kly sovet, Ah SSSA (Astrono	midal opinale, me o	oùh,
	20 mr6.	ENGL: 00	OF OIL I	DC.ES
	1: 005	OTHER: CCC		

ACC NR. 416014954

SOURCE CODE: UR/0227/65/000/008/0023/0025

AUTHOR: Yeliseyev, Yu. A.; Voroshilin, Ye. A.; Biyevets, N. L.; Krylov, A. G. 24

B

ORG: none

TITIE: Construction of a container glassware storage warehouse of reinforced concrete

SOURCE: Promyshlennoye stroitel'stvo, no. 8, 1965, 23-25

TOPIC TAGS: reinforced concrete, construction, lacquer, corrosion protection

ABSTRACT: A description is given of the construction of a 24 X 48 meter warehouse with supporting frame made of prefab arches each consisting of six straight sections of reinforced concrete, bolted together. The prefab sections were compacted, heat-hardened for 4 hours at 70°C, reinforced with steel mesh and given an anti-corrosion coating of bituminous lacquer. They were then stored in special holding racks, in which they were also transported to the construction. Photographs show the forming, transporting and assembly of the individual straight sections into arches, as well as the completed warehouse. A table shows the expenditure of materials manpower and money per square meter of horizontal projection involved in the construction. Orig. art. has: 5 figures and 1 table. [JPRS]

SUB CODE: 13 / SUBM DATE: none

Card 1/1 BK

UDC: 624,023.8:725.35

APPROVED FOR RELEASE: 06/14/2000

CIA-RDP86-00513R000826820012-2"

USER / Forest Science. Biology and Typology of Trees.

K-2

Abs Jour

: Ref. Zhur - Biologiya, No 17, 1958, No. 77487

Author

: Rrylov, A. G.

Inst

: West Siburian Branch, AS USER

Titlo

: On Types of Dark Conifer Forests in the Biyn River Basin

Orig Pub

: Tr. po lesn. kh-vu Zap. Sibiri. Zap.-Sib. fil AN SSSR,

1957, vyp. 3, 167-174

Abatract

: Fir and codar types of forest developed in the summer of 1956 in the Biya River Basin are briefly characterized. In the group of shortgrass variety firs, there were noted form-sedge, sanicle-wood sorrel poplar fir. In the group of mixed firs, spiraea-bracken-herbaceous fir were noted. In the codar forests, there were isolated: cedar-whortleborry, herbaceous-messy codar, wood sorrel-messy codar (group of messy cedars); form-sedge cedar (group of grass variety cedars); form-beach grass cedar and herbaceous

Card 1/2

TITLE THE TALL

Moscow Automobile Plant at the 40th anniversary of the October

Revolution. Avt.1 trakt.prom. no.10:8-13 0 '57. (MIRA 10:12)

1. Moskovskiy avtosavod imeni Likhacheva.
(Moscow---Automobile industry)

KRYLOV, A.G.

Automatic machines in shops of the Likhachev Automobile Plant. Za rul. 17 no.8:3-5 Ag '59. (MIRA 12:12)

1. Direktor avtomobil'nogo zavoda im. I.A.Likhacheva.
(Moscow--Automobile industry)
(Automation)

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

CHUMAKOV, N.M., red.; KIREYEV, M.I., red.; AKULOV, Ye.F., red.; IVANOV, N.N., red.; KNYAZEV, P.I., red.; CHICHILO, I.K., red.; KRYLOV, A.G., red.; GLUSHKO, G.I., tekhn. red.


[Safety engineering regulations for operating and servicing electrical systems of industrial enterprises required for the industrial plants of economic councils, ministries, and departments]Pravila tekhnicheskoi ekspluatatsii i bezopasnosti obsluzhivaniia elektroustanovok promyshlennykh predpriiatii; obiazatel'ny dlia promyshlennykh predpriiatii sovmarkhozov, ministerstv i vedomstv. Utverzhdeny 10 fevralia 1961 g. Moskva, Dnepropetrovskoe knizhnoe izd-vo, 1962. 279 p.

(MIRA 16:3)

1. Russia (1923- U.S.S.R.) Glavnoye energeticheskoye upravleniye.

(Electric power distribution-Safety regulations)

ERYLOV, A.1. New type of roller feed. Kuz.-shtam. proizv. 3 no.11:41-42 H (Feed mechanisms) (Forging machinery) 



KRYLOV, A.I., insh.

Modernizing the engaging friction clutch. Vest.msh. 41 no.7:88
Jl '61. (Clutches (Machinery))

(Clutches (Machinery))

L 19638-63 EPF(c)/EWP(q)/EWT(m)/BDS AFFTC/ASD Pr-4 JD S/0056/63/045/003/0464/0468

AUTHORS: Zherebin, Ye. A.; Krylov, A. I.; Polikarpov, V. I.; Yuzvuk, N. N.

---

TITLE: Investigation of the gamma radiation from Cs-140

SOURCE: Zh. eksper. i teoret. fiziki, v. 45, no. 3, 1963, 464-468

TOPIC TAGS: Cs-140, gamma radiation, short-lived fragment, spectral line

ABSTRACT: A method for investigating the gamma rays from the short-lived (half-life 66 sec) fragment Cs<sup>140</sup> is described, along with the gas loop used to supply the Xe and Kr to the measurement place and to enrich the mixture of the decay product with the investigated fragment product. The Cs<sup>140</sup> was investigated by a high-speed chemical separation of the cesium. The lines 0.59 ± 0.01, 0.88, 1.14, 1.62, 1.85, 2.06, 2.32, 2.72, 3.15 MeV were observed as a result in

Card 1/82/

L 19638-63

ACCESSION NR: AP3007063

6

the gamma rays. "In conclusion, the authors thank Ye. A. Tamonov and O. V. Chubakov for useful discussions and advice, and also A. N. Draskov, A. G. Dudoladov, Ye. A. Gershanov, and A. V. Morozov for directly participating in the experiments." Orig. art. has 6 figures.

ASSOCIATION: None

SUBMITTED: 29Mar63

DATE ACQ: 080ct63


ENCL: 01

SUB CODE: PH

NO REF SOV: 001

OTHER: 002

Card 2/82



PRAGA, D. T., Eng.; KRYLOV. A. I.; GLUSHAYENKO, E. I.

Knitting Machines

Single-process method of making socks with flat knitting machines. Leg. prom. 12 No. 9, 1952.

1952 9. Monthly List of Russian Accessions, Library of Congress, December \_109758, Uncl.

KHOLODOV, Vladimir Ivanovich; ERYLOV, Aleksandr Ivanovich; TEPIMAKHOVA,
M. Ta., red.; LECHOVA, L.F., teknin.red.

[Through the years (1858-1958)] Skvos' gody, 1858-1958 gg.
Vladimir, Vladimirakoe knizhnoe izd-vo, 1958, 97 p. (MIRA 12:9)

(Sobinka--Textile workers)

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

KOZENKO, K.M., inzh.; KRYLOV, A.I., inzh.; BELOUS, N.G., inzh. New techniques used in manufacturing warp-knitted artificial furs.

Isv.vys.ucheb.zav.; tekh.leg.pron. no.6:75-85 158. (MIRA 12:4)

1. TSentralinaya nauchno-eksperimentalinaya laboratoriya trikotashnoy promyshlennosti Gosplana USSR. (Fur, Artificial)

CIA-RDP86-00513R000826820012-2" **APPROVED FOR RELEASE: 06/14/2000** 

下口也和阿拉拉克的医别的人物的 网络克拉斯 计数字记录符 KOZENKO, K.H.; KRYLOV, A.I.; BELOUS, N.G. Developing the technology of knitting pile fabrics for artificial fur. Tekst.prom. 19 no.1:44-47 Ja '59. (HIRA 12:1) (Knitting, Machine) (Fur, Artificial)

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

SMIRNOV, Leonid Stepanovich; GONTARENKO, Aleksandr Nikolayevich;
GORDIYENKO, Mariya Georgiyevna; KRYLOV, Aleksandr Iosifovich;
NOVAK, Nikolay Stepanovich; INASHCHRIKO, T.V., red.; STARDUB,
T.A., tekhn. red.

[Manufacture of artificial fur] Proizvodstvo iskusstvennogo
mekha. Kiev, Gos. izd-vo tekhn. lit-ry USSR, 1961. 138 p.

(Artificial fur)

(Artificial fur)

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

# "APPROVED FOR RELEASE: 06/14/2000


CIA-RDP86-00513R000826820012-2

KRYLOV, A.I. [Krylov, O.I.]; SKIBA, L.A.

Economic efficiency of the manufacture and the of nylon yarn produced with a simplified method. Leh. prom. no.A:66-69
C-D '65. (MIRA 19:1)

### "APPROVED FOR RELEASE: 06/14/2000


### CIA-RDP86-00513R000826820012-2



# "APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2

the bottom of the latter (see Fig. 1.). Orig. art. has: 1, figure. [AZ]  SUB CODE: 11,13/SUBM DATE: 29Jan65/ ATD PRESS: 5-0//							
	,						
				-			
			44		· -"		
		•		e.			
					•		
Card 2/2 (¹(',							

```
The KEK-1,0 combine for bean harvesting. Biul.tekh.-ekon.inform.
no.2:55-56 '58. (MIRA 11:4)
(Combines (Agricultural machinery))
(Beans--Harvesting)
```



## "APPROVED FOR RELEASE: 06/14/2000

CIA-RDP86-00513R000826820012-2

KRITADV, A.I.; FROIPVOLOVA, G.A.

Numerical study of liquid flow between rotating cylinders. Sbor. rab. VTS 1980 2:174-181 163. (1984 17:7)

### "APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2

KRYLOV, A.L.; MIGNIK, A.F.

Stability of viscous flow between two rotating cylinders. Sbor. rab. VTS MGU 2:182-189 '63. (MIRA 17:7)

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

AUTHOR:

Krylov, A.L.

20-1-10/54

TITLE:

The Boundary Value Problem for the Nonlinear Equation of Second Order in a Bounded Domain with a Degenerate Boundary. (Pervaya Krayevaya zadacha dlya nelineynogo ellipticheskogo uravneniya vtorogo poryadka ogranichennoy oblasti s vyrozhdennoy granitsey)

PERIODICAL: Doklady Akademii Nauk SSSR, 1957, Vol 115, Nr 1, pp.42-44 (USSR)

ABSTRACT:

The present paper treats this problem for the Lagrange equation of a certain variation problem. In contrast to the linear case the boundary values in the case of a very large class of nonlinear problems may be assumed on varieties with any dimension number, provided that this dimension number is smaller than the dimension number of the initial space. E.g. the boundary in the two-dimensional case may consist of a closed line bounding the domain and of a finite number of smooth lines within this domain. The obtained solutions are continuous in such a domain and in the case of smooth integrands they are solutions in the classical sense. Reference is made to some relevant earlier works. The author here examines the following example: K signifies the circle of unit radius on a plane with a cut-out center in the origin of coordinates and

Card 1/3

The Boundary Value Problem for the Nonlinear Equation of Second 20-1-10/% Order in a Bounded Domain with a Degenerate Boundary.

I(u) =  $\iint \{\alpha(u_x^2 + u_y^2)^2 + \beta(u_x^2 + u_y^2)\}$  dxdy applies. The function  $u_0 \in \mathbb{V}_4^{(1)}$  is sought which on the boundary K assumes the values  $\mathfrak{P}(Y)$  when  $x^2 + y^2 = 1$ ,  $\mathfrak{P}_0$  when  $x^2 + y^2 = 0$ . In this connection  $\mathfrak{P}(Y)$  is attainable 'allowable' for  $\mathbb{V}_4^{(1)}$ . When  $\alpha = 0$ , this problem has a solution. When  $\alpha > 0$ ,  $\alpha > 0$  the problem is regular, i.e. in the case of a smooth solution it is reduced to a non-degenerate elliptical equation. In the case of  $\alpha = 0$  the solution  $\alpha = 0$ .

A solution u (x,y) of the variation problem is determined here which takes the assumed values on the boundary of K and this u (x,y) is a unique solution of the variation problem smooth in K. A relevant theorem is given. This theorem can be generalized to the case that F(in the above-mentioned formula) depends on independent variables and functions, and it permits a considerable weakening of the other conditions. There are 6 Slavic references.

Card 2/3

The Boundary Value Problem for the Nonlinear Elliptic Equation 20-1-10/54 of Second Order in a Bounded Domain with a Degenerate Boundary.

ASSOCIATION: Moscow State University imeni M.V.Lomonosov (Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova)

PRESENTED: January 28, 1957 by S.L. Sobolev, Academician

SUBMITTED: January 26, 1957.

AVAILABLE: Library of Congress

Card 3/3

problem for certain quasi-linear equations in the appear with degenerated boundary." Mos, 1958, 8 pp (Mos State Univim M.V. Lomonosov. Mechanica Mathematics faculty) 150 cories. Pibliography at end of text (11 titles) (KL, 27-58, 102)

- 15 -

AUTHOR:

Krylov, A. L.

20-118-5-9/59

TITLE:

Propagation of Limit Equilibrium in the Axially-Symmetrical Two-Dimensional Elastic-Loose Body Case (Osesimmetricheskaya ploskaya uprugo-sypuchaya zadacha rasprostraneniya sostoyaniya predel'nogo

ravnovesiya)

PERIODICAL:

Doklady Akademii Nauk SSSR, 1958, Vol. 118, Nr 5,

pp. 882-883 (USSR)

ABSTRACT:

§ 1: The setting of the problem and the hypothesis of motion: The author here investigates the two-dimensional deformed state of a medium which fills the external part of the circle of unit radius on the x,y-plane. In the case of low stresses the medium is described by the equations of linear elasticity theory, in the case of greater stresses, however, the medium adopts loose properties. Starting at the moment t = 0 the sector begins to swell according to the law  $Y = Y_1(t), Y_1(0) = 1$ , and to rotate according to the law

Card 1/4

Propagation of Limit Equilibrium in the Axially-Symmetrical 20-118-5-9/59 Two-Dimensional Elastic-Loose Body Case

the surroundings of the sector. The stress and the displacement in the elastic and in the loose zone are to be determined as well as of the boundary  $\gamma(t)$  between these zones, furthermore the admissibility of the boundary conditions  $\{r_1(t), \psi_1(t)\}$  for the maintainance

of the equilibrium on the boundary is to be investigated. Two hypotheses of motion were used for the solution of this problems: the hypothesis by Mises (Mizes), and a hypothesis by the author which runs as follows: The agreement of the direction of the maximum displacement velocity with one direction of the slide lines (i.e. with those forming the smallest angle with the limit on the boundary) is suggested, at the same time maintaining the incompressibility condition. The problem without rotation is instable in the hypothesis discussed here; in the case of presence of a small rotation the movement differs considerably from mere swelling. § 2 solves the problem with swelling explicity for both hypotheses. The solution of the problem by means of the hypothesis of Mises (Mizes) leads to a paradoxical conclusion on the unlimited capability of loosening of the

Card 2/4

Propagation of Limit Equilibrium in the Axially-Symmetrical 20-118-5-9/59 Two-Dimensional Elastic-Loose Body Case

> medium for great radii of the sector. § 3 gives the equations of equilibrium which are to be integrated in the solution of the problem with rotation in the loose zone. In the elastic zone the solution is described by formulae by S. Sobolev (Ref 1). Then a differential equation for the determination of the displacement is given on the condition of the author's hypothesis. If the stability of the stress tensor, the displacements, and of the velocities is used for  $\gamma$  (t), a closed system of equations is obtained for the determination of the unknown time functions. All unknown functions are finally expressed by the solution A (t) of the Cauchy problem for an integral-differential equation given here. § 4 investigates quite briefly the admissibility of boundary movement. In the rotation of the boundary with given velocity the slide lines touch the boundary. In the case of increasing velocity the boundary of the medium ruptures along the slide lines. There are 4 references, 4 of which are Soviet

Card 3/4

### "APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2

Propagation of Limit Equilibrium in the Axially-Symmetrical Two-Dimensional Elastic-Loose Body Case

20-118-5-9/59

14

PRESENTED:

September 5, 1957, by S. L. Sobolev, Member, Academy of

Sciences USSR

SUBMITTED:

December 24, 1955

Card 4/4

AUTHOR:

Krylov, A.L.

20-119-5-6/59

TITLE:

Boundary Value Problems and Biorthogonal Decompositions in Banach Spaces (Krayevyye zadachi i biortogonal'nyye razlozheniya v

banakhovykh prostranstvakh)
SSSR

PERIODICAL: Doklady Akademii Nauk, 1958, Vol 119, Nr 5, pp 865-867 (USSR)

ABSTRACT:

With the aid of the theorem of Calderon and Zygmund [Ref 3] on the conditions for the boundedness of the singular integral operator J:

 $Jf \equiv \int k(P,Q)f(Q)dQ$ 

the author succeeds in extending essentially the results of Vishik [Ref 2] basing on the use of the orthogonal projections. While Vishik at first obtained a biorthogonal decomposition of the considered space on a geometric way and then applied it to the solution of boundary value problems, the author at first solves the boundary value problem with the aid of the theorem of Calderon-Zygmund, wherefrom then there follows the decomposition. The representation is made with the example of the Laplace equation, but it remains true for equations of higher order and also for systems with smooth coefficients.

Card 1/2

There are 7 references, 5 of which are Soviet, 2 American.

#### "APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2

Boundary Value Problems and Biorthogonal Decompositions in Banach 20-119-5-6/59 Spaces

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im.M.V. Lomonosova (Moscow State University im.M.V.Lomonosov)

PRESENTED: November 23, 1957, by S.L.Sobolev, Academician SUBMITTED: November 22, 1957

Card 2/2

CIA-RDP86-00513R000826820012-2" APPROVED FOR RELEASE: 06/14/2000

AUTHOR:

Krylov L.L.

SOV/20-121-5-7/50

TITLE:

On a Necessary and Sufficient Condition that a Function Belong to the Class W(1) of Sobolev (Ob odnom neobkhodimom i dostatochnom priznake prinadlezhnosti funktsii klassu Wp Soboleva)

PERIODICAL: Doklady Akademii nauk SSSE, 1958, Vol 121, Nr 5, pp 795-796 (USSR)

ABSTRACT:

In the domain  $\Omega$  with the boundary ! let the imbedding theorems [Ref 3] be valid (e.g. let [ be smooth). Let \(\Omega\) be subdivided by the straight lines x = x<sub>1</sub>, 1=0,1,..., m-1, y = y<sub>1</sub>, j=0,1,...n-1; let & be the smallest distance between the parallel straight lines of the subdivision. Let a continuous function F(x,y),  $(x,y) \in \Omega$ be of bounded p-variation (p > 2) if the sums

$$\sum_{i,j} \frac{|P(x_i + h_i y_j) - P(x_i y_j)|^{\frac{p}{P}}}{h^{p-2}}, \sum_{i,j} \frac{|P(x_i y_j + h) - P(x_i y_j)|^{\frac{p}{P}}}{h^{p-2}}$$

are bounded from above by a number v not depending on the subdivision and on  $h(h \le \beta/2)$ . It is summed over the points  $(x_1, y_1)$  lying in  $\Omega$  with its  $\beta$ -neighborhood. Theorem: In order that a function belongs to the class  $\mathbf{w}_{n}^{(1)}$ 

Card 1/2

On a Necessary and Sufficient Couldition that a Fraction Belong 300/20-121-5-7/50 to the Class  $\mathbb{W}_p^{(1)}$  of Sobolev

it is necessary and sufficient that it is of bounded p-variation. There are 3 references, 2 of which are Soviet, and 1 Hungarian.

ASSOCIATION: Moskovskiy gosudarstvennyy universiteb imeni M.V. Lomonosova (Moscow State University imeni M.V. Lomonosov)

PRESENTED: April 19, 1958, by S. L. Sobolev, Academician

SUBMITTED: April 7, 1958

Card 2/2

16(1) AUTHOR:

Krylov, A.L.

SOV/20-124-2-5/71

TITLE:

First Boundary Value Problem for Some Quasilinear Equations of Parabolic Type in the Banach Space (Pervaya krayevaya zadacha dlya nekotorykh kvazilineynykh uravneniy parabolicheskogo tipa v banakhovom prostranstve)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 124, Nr 2, pp 264-267 (USSR)

ABSTRACT:

Let  $\Omega$  be a bounded domain of the maximum dimensional space  $\mathbf{x}_{-}(\mathbf{x}_{1},\ldots,\mathbf{x}_{m})$ with the boundary Consisting of manifolds of m-1,m-2,...,1,0 dimensions. Let  $Q = \Omega \times [0,T]$  be a cylinder in the space (x,t). In Q the author considers the equation

 $u_{t} - \sum_{i=1}^{m} \frac{\partial}{\partial x_{i}} (F_{u_{x_{i}}}(u_{x_{i}}, \dots, u_{x_{m}})) = f(x, t)$ (1)

with the boundary conditions

$$\frac{u}{|s|} = 0 \qquad u|_{t=0} = \varphi(x),$$

where s=  $\Gamma \times [0,T]$ . It is assumed that for  $\Omega$  and  $\Gamma$  there hold the imbedding theorems of Sobolev, that for a restriction to smooth

Card 1/2

u(x,t) the functionals  $\int_{\Omega} Fdx$  and  $\int_{\Omega} F dxdt$  are uniformly convex,

APPROVED FOR RELEASE: 06/14/2000

CIA-RDP86-00513R000826820012-2"

First Boundary Value Problem for Some Quasilinear SOV/20-124-2-5/71 Equations of Parabolic Type in the Banach Space

and that  $\sum_{\mathbf{x_i}} \mathbf{x_k} \mathbf{x_i} \mathbf{x_k} \leq \mathbf{x} \leq \mathbf{x} \leq \mathbf{x} \leq \mathbf{x}$ ,  $\mathbf{x} > 0$ . The author proves

the existence of a unique generalized solution of (1). The proof is made according to the method of Galerkin supported by considerations of variation.

There are 3 Soviet references.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova (Moscow State University imeni M.V. Lomonosov)

PRESENTED: September 1, 1958, by S.L.Sobolev, Academician

SUBMITTED: April 17, 1958

Card 2/2

· 1. 年度中,1860年的開發中格克拉爾聯盟開展的開始。至底於1. 年初了一年

ZHIDKOV, N.P.; KORNEYCHUK, A.A.; KRYLOV, A.L.; MOSTINSKAYA, S.B.

Plane-parallel motion of a viscous liquid between two rotating cylinders. Vych. met. i prog. 1:152-166 '62. (MIRA 15:8) (Fluid dynamics)

KRYLOV, A.L.

Models involving a finite number of degrees of freedom for a certain class of problems in mathematical physics. (Difference systems involving a law of conservation). Dokl. Ad SSSR 142 no.3:572-575 Ja 62. (NIRA 15:1)

Moskovskiy gosudarstvennyy universitet im. E.V. Lomonosova.
 Predstavleno akademikom S.L. Sobolevym.
 (Mathematical physics) (Difference equations)

s/020/62/146/001/005/016

AUTHOR:

Krylov, A. L.

TITLE:

Argument establishing Dirichlet's principle for the first boundary value problem in the non-linear theory of elasticity

PERIODICAL: Akademiya nauk SSSR. Doklady, v. 146, no. 1, 1962, 54 - 57

TEXT: The problem is to find a displacement vector u(x) which agrees with a given vector  $\phi$  at the boundary  $D\Omega$  of the domain  $\Omega$  and which corresponds to the minimum value of the integral

 $g(\beta)d\beta$ ) $d\Omega$ , where k is a constant,  $\epsilon$  and  $\Gamma^2$  are the linear and square invariants of the deformation tensor | Eik and of the deviation tensor  $\|\xi_{ik} - \delta_{ik} \xi\|$ , respectively, and where the function  $g(\Gamma^2)$ 

is defined by the fundamental relations of the non-linear theory of elasticity in the following way:  $D_{\xi} = \psi D_{\sigma}$ ,  $\psi = f(T)$ ,  $T = 2g(\Gamma^2)\Gamma$ . components of the vector u are connected with those of the deformation Card 1/2

Argument establishing Dirichlet's...

s/020/62/146/001/005/016 B112/B108

tensor by the usual relation  $\varepsilon_{ik} = (1/2)(\partial u_i/\partial x_k + \partial u_k/\partial x_i)$ . The solvability of this problem is shown by means of a solving requence  $u^{n} \in W_{\beta}^{(1)}$ , where  $\beta$  is defined by the representation  $g(\hat{\Gamma}^{2}) = B(\hat{\Gamma}^{2})(\beta-2)/2$ .

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University imeni M. V. Lomonosov)

PRESENTED:

March 28, 1962, by S. L. Sobolev, Academician

SUBMITTED:

March 15, 1962

Card 2/2

**APPROVED FOR RELEASE: 06/14/2000** CIA-RDP86-00513R000826820012-2"

3/794/62/000/001/006/010

AUTHORS: Zhidkov, N.P., Korneychuk, A.A., Krylov, A.L., Mastinskaya,

The plane-parallel motion of a viscous fluid between rotating cylinders. TITLE:

Vychislitel'nyye metody i programmirovaniye; sbornik rabot SOURCE:

Vychislitelinogo tsentra Moskovskogo universiteta. no. 1. Ed. by N.P. Trifonov, G.S. Roslyakov, and Ye. A. Zhogolev. [Moscow] Itd-vo

Mosk. un-ta, 1962, 152-166.

The paper is intended to investigate the motion of a fluid by means of the direct numerical integration of the nonlinear Navier-Stokes equations. The direct objective of the investigation is the case of a viscous incompressible fluid in an x, y, z space contained between two infinite cylinders having radii  $R_1^2 \le x^2 + y^2 \le R_2^2$ which rotate with different angular velocities. The behavior of the fluid in a plane solution (i.e., independent of z) is to be found. Upon establishment of the necessary matrix expression the solution of the problem on the machine "Strela" at the Computing Center of the Moscow State University is described. The program consists of 4 separate parts, (1) the shaping and the memory formulation of the distribution program, (2) the program of the calculating of coefficients, (3) the program of preparation of the initial data, and (4) the program of the solution of the difference equations.

Card 1/2

The plane-parallel motion of a viscous fluid ....

5/794/62/000/001/006/010

Calculations were made for 4 Reynolds numbers, namely, 200, 500, 1,000, and 2,000, with different timewise steps. It was found that in the largest network employed the eddies smoothened out and the flow became smooth and trivial. The time of establishment grew proportional to the R number. In all probability this network was exceedingly crude and had its own viscosity. The use of a fine network revealed a turbulent motion, if the degree of turbulence is intended to signify the number of eddies in the flow. The amplitude of the perturbations decreases with time, but turbulence grows and new eddies are formed, even though with smaller amplitudes. Ultimately, the motion in the network becomes smooth, since the network does not admit eddies of a size smaller than the dimensions of the network. Future work will comprise a reduction of the size of the network by one-half. Thanks are expressed to I. M. Gel'fond, corresponding member, AS USSR, for overall direction of the work and many valuable specific advices. There are 4 figures and 1 Russian-

Card 2/2

\$/3043/63/000/002/0165/0173

AUTHOR: Kry\*lov, A. L.

TITLE: Finite-dimensional models for equations of mathematical physics

SOURCE: Hoscow, Universitet. Vy\*chisliteliny\*y tsentr. Sbornik rabot, no. 2, 1963. Chislenny\*ye metody\* v gazovov dinamike, 165-173

TOPIC TAGS: finite dimensional model, energy conservation law, Cauchy problem, viscous incompressible finid model periodic boundary condition, mathematical physics equation  $\frac{\partial}{\partial t} \left( \sum_{i} q_i L^i q_i - L^0 \right) + \sum_{i} \frac{\partial}{\partial t} \left( \sum_{i} q_i L^i q_i - L^0 \right) = 0$ 

ABSTRACT: A broad class of partial differential equations of mathematical physics possesses the following property: the equations themselves express a law of conservation of certain physical quantities, and, in addition, a supplementary law of conservation arises as a consequence of the realization of the fundamental system (e.g., conservation of energy in crystal optics). Following the work of S. K. Godunov the author examines equations of the type

$$\frac{\partial L^{0}q_{i}}{\partial t} + \sum_{l=1}^{n} \frac{\partial L^{l}q_{i}}{\partial x_{l}} = 0, \quad l = 1, 2, \dots, m,$$

$$L^{k} = L^{k}(q_{1}, q_{2}, \dots, q_{n}), \quad k = 0, 1, \dots, n.$$
(1)

Card 1/4

The equations of crystal optics and others can be carried over to this form. The law of conservation fulfilled in the solutions of (1) is

(2)

the law of conservation of entropy or energy or, for the irreversible process, the dissipation of energy. The purpose of the present paper is to obtain finite dimensional models for such systems having "point" law energy conservation of type (2). A. N. Kolmogorov first dealt with this, regarding turbulence, for Navier-Stokes equations of a viscous incompressible liquid. It is possible that study of the ergodic preparties of such a model would lead to conclusions for the laminar case, although the author knows of no such work. He first lets [=j=1], which gives

$$\frac{\partial}{\partial t} A_{q}(q) + \frac{\partial}{\partial x} B_{q}(q) = \frac{\partial}{\partial x} \left( C(q) \frac{\partial q}{\partial x} \right)$$
(3)

(on the right is the dissipation member). For simplicity he examines the problem of Cauchy with periodicity in x:

$$\frac{\partial^{k} q}{\partial x^{k}}(l,0) = \frac{\partial^{k} q}{\partial x^{k}}(l,l), \qquad k = 0; 1$$
 (4)

Card 2/4

Similarly, it should be possible to examine other boundary conditions and the case of an unrestricted x decreasing towards infinity. Transforming (3) and utilizing (4), an identity is established (the law of dissipation of energy). A finite-dimensional model of problem (3), (4), is constructed to permit similar transformations on a difference equation approximating (3). This is done by dividing the interval of periodicity into uniform segments and defining the desired functions  $q^k$ , their derivatives, and the derivatives of their functions at the division points. The transformation of equation (3) by approximation is described in detail.

 $\frac{\partial L^{0}q_{i}}{\partial t} + \sum_{\ell=1}^{n} \frac{\partial U_{q_{\ell}}}{\partial x_{j}} = \sum_{l,l,s} \frac{\partial}{\partial x_{j}} b_{li}^{ls} \frac{\partial q_{l}}{\partial x_{s}}, \qquad l = 1, 2, \dots, m$ (5)

with symmetry and positively defined matrices, the constructions are carried through in the same fashion. Unfortunately, for more than one space variable, only a tetrahedral grid can be established. With reasoning similar to this, V. I. Lebedev was the first to use such a grid for the same goal. For simplicity, two space variables are considered. The differential operators are defined and discussed and the establishment of the grid system is described in detail. The results of applying this to equation (5) are given. Finally, the author examines the solution of the Navier-Stokes equation for viscous incompressible liquids.

Card

ing down difference schemes with the same properties. He points out that, in the works of A. N. Tikhonov and A. A. Samarskiy, such schemes are constructed and studied for a very broad class of ordinary differential equations. Orig. art. has: 28 formulas.

ASSOCIATION: Vy\*chislitel\*ny\*y tsentr MGU (Computer center, Moscow State University)

SUBMITTED: 00

DATE ACQ: 16Dec63

ENCL: 00

SUB CODE: AI

4/4

NO REF SOV: 003

OTHER: 000

Card

APPROVED FOR RELEASE: 06/14/2000

CIA-RDP86-00513R000826820012-2"

\$/3043/63/000/002/0174/0181

AUTHOR: Kry\*lov, A. L.; Proizvolova, Ye. K.

TITLE: Numerical study of flow between two rotating cylinders

SOURCE: Moscow. Universitet. Vy\*chisiitei\*ny\*y tsentr. Sbornik rabot, no. 2, 1963, Chislenny\*ye metody\* v gazovoy dinamike. 174-181

TOPIC TAGS: viscous incompressible flow, flow between rotating cylinders, flow equation numerical solution, finite difference approximation, successive approximation method, flow stability, axisymmetric flow, viscous flow, incompressible fluid, viscous incompressible fluid, viscous flow, incompressible flow, curved flow, recurrence formula

ABSTRACT: The authors have calculated the behavior of a streamlined flow of a viscous incompressible fluid between coaxial cylinders of infinite height, rotating at angular velocities  $\Omega_1$  and  $\Omega_2$ , in relation to definite relationships between Reynold's numbers Rej and Reg. The flow is assumed to be periodic in relation to height and asymmetric. The phenomena discussed occur within the Reynold's number range of 100 to 200. The problem was postulated in such a manner that the Reynold's number was governed by the non-dimensional viscosity factor V. The authors attempted to clarify the periodic patterns possible at a fixed Reynold's 1/62.

Card

number ( $\forall$ = 1/200) and, particularly, to establish the presence of more intensive frequencies in a spectrum component which is unstable with respect to infinitely small turbulences. Calculations were made for the variants:

$$y = \frac{1}{200}$$
;  $H = l, 2l, 4l, \left(l = \frac{\pi}{3.2}\right)$ 

Results are illustrated by several graphs (see Figs. 1, 2, 3 and 4 in the Enclosure). "In conclusion, we express our gratitude to I. M. Gel'fand, whose influence and participation were evident in originating theoretical solutions to problems on the stability of flow of viscous fluids, as well as to S. B. Mostinskaya and Yu. Gormatyuk for their participation in programming." Orig. art. has: 4 graphs and 9 formulas.

ASSOCIATION: Vy\*chislitel'ny\*y tsentr MGU (Computer center, Moscow State University)

SUBMITTED: 00

DATE ACQ: 16Dec63

ENCL: 04

SUB CODE: AI

NO REF SOV: 003

OTHER: 001

2/6 >

Card

\$/3043/63/000/002/0182/0189

ACCESSION NR: AT4006717

AUTHOR: Kryklov, A. L.; Hisnik, A. F.

TITLE: Stability of viscous flow between two rotating cylinders

SOURCE: Moscow. Universitet. Vy\*chisliteliny\*y tsentr. Sbornik rabot, no. 2, 1963. Chislenny\*ye metody\* v gazovoy dinamike, 182-189

TOPIC TAGS: viscous flow stability, flow between rotating cylinders, Taylor flow equation, viscous incompressible flow, finite difference method, neutral curve equation, flow equation integration, viscous fluid, incompressible fluid, viscous Incompressible fluid, flow stability equation, flow equation spectral property, viscous flow, incompressible flow, Taylor flow, curved flow stability

ABSTRACT: The article presents the results of numerical integration of the Orr-Sommerfeld equations for Taylor's flow. This concerns a flow of incompressible fluid between concentrically rotating cylinders of infinite height. The concepts of critical Reynold's number and neutral curve are defined. The latter was plotted In a plane  $(R, \mathcal{R})$ , where R is the Reynold's number of a streamline flow and is the frequency of stability loss, while turbulence is written as f(r)ect+ iz (see graph in the Enclosure). Calculations were carried out on a "Strela" computer at the Vy\*chislitel'ny\*y tsentr HGU (Computer center of Hoscow State Cord 1/3 Card

University). It was found that the equation for stability of a Taylor flow exhibits spectral characteristics which differ strongly from those of the corresponding equation for a Poiseuille flow. Orig. art. has: I table, I graph, 6 formulas.

ASSOCIATION: Vy\*chisliteliny\*y tsentr MGU (Computer center, Moscow State

University)

SUBMITTED: 00

DATE ACQ: 16Dec63


ENCL: 01

SUB CODE: AI

NO REF SOV: 000

OTHER: 005

Card 2/3



```
s/020/63/148/001/001/032
B172/B186
```

Uniform distribution of ...

s/020/63/148/001/001/032

where dense on  $S^2$ , then their distribution on  $S^2$  is uniform. This theorem may be understood as an ergodic hypothesis in which a semi-group with two generatrices plays the role of time. The authors generalize this theorem; S2 is replaced by a homogeneous space on which a compact Lie group is defined, whereas a free group with two generatrices or a group with three generatrices a,b,c, for which abc-e is valid, is taken as "time". Then, sets of linear differential equations

= A(z)xare considered

 $(z = a complex variable, x = a vector <math>(x_1, ..., x_n)$  of the n-dimensional complex space, A = a matrix which, except for three singular points lying on a Riemann sphere, depends analytically on z). The following theorem is formulated: If the monodromic group of such a system is bounded, then the system has a unique first integral  $(B(z)x,\overline{x}) = const$ , where B(z) is a positive definite, self-adjoint matrix. For Gauss' hypergeometric equation, the assumption of this theorem is fulfilled. Arising from this, the authors mention finally a number of unsolved problems.

Card 2/3

Uniform distribution of ... S/020/63/148/001/001/032 B172/B186 ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University imeni M. V. Lomonosov)

PRESENTED: June 23, 1962, by A. N. Kolmogorov, Academician

SUBMITTED: June 18, 1962

Card 3/3

# KRYLOV, A.L.

Proof of the instability of a certain flow of a viscous incompressible fluid. Dokl. AN SSSR 153 no.4:787-789 D 163. (MIRA 17:1)

1. Moskovskiy gosularstvennyy universitet im. M.V. Lomonosova. Predstavleno akademikom A.N. Kolmogorovym.

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

KRYLOW, A.L. (Moscow)

"Some of problems stability of viscous fluid flow".

report presented at the 2nd All-Union Congress on Theoretical and Applied Mechanics, Moscow, 29 Jan - Feb 64.

#### KRYLOV, A.L.

Stability of Poiseuille flow in a plane channel. Pokl. AN SSSR 159 no.52978-981 11 164 (MIRA 1821)

1. Moskovskiy gosudarstvennyy universitet. Predstavleno akademikum A.N. Kolmogorovym

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

L 00362-66 EWT(1)/EWP(m)/EWA(d)/FCS(k)/EWA(1)

ACCESSION NR: AT5013285

UR/3043/65/000/004/0108/0114

AUTHOR: Krylov, A. L.

TITLE: Proof of the instability of Kuetta flow between rotating cylinders

SOURCE: Moscow. Universitet. Vychislitel'nyy tsentr. Sbornik rabot, no. 4, 1965. Chislennyye metody v gazovoy dinamike (Numerical methods in gas dynamics), 108-114

TOPIC TAGS: Kuetta flow, laminar flow, cylindric flow, flow analysis

ABSTRACT: The flow of a viscous incompressible fluid between rotating coaxial cylinders is one of the most intensively and successfully studied problems concerning cases of loss of stability of laminar stationary flow and a subsequent transition to another laminar stationary flow. The problem concerning the stability of the laminar Kuetta flow reduces to the question of the existence in the system

with the boundary condition

$$(L-\lambda^2-\sigma R) = 2\lambda^2 R\omega v,$$

$$(L-\lambda^2-\sigma R) v = 2\lambda^2 R\omega v,$$

$$u = v = \frac{du}{dr} = 0 \qquad r = R_1(l = 1, 2)$$

Card 1/2

L 00362-66

ACCESSION NR: AT5013285

of an eigenvalue  $\zeta$  (R,  $\chi$ ) in the right half-plane Re  $\zeta$  > 0. Here

 $L = \frac{d}{dr} \frac{1}{r} \frac{d}{dr} r, \quad R_1 < r < R_2$ 

and R > 0 is the Reynolds number; l > 0. Taylor solved this problem numerically in 1923. The present paper brings a mathematically rigorous solution to the problem. Orig. art. has: 22 formulas.

ASSOCIATION: Vychislitel'nyy tsentr, Moskovskiy universitet (Computer Center, Moscow University)

SUBMITTED: 00


ENCL: 00

SUB CODE: ME, MA

NO REF SOV: 006

OTHER: 002

Card 2/2



Over-all mechanisation in tempering shops. Mashinostroitel'
no.10:1-5 0 '57. (MIRA 10:11)
(Automatic control) (Tempering) (Sverdlovsk--Bearing industry)

KRTLOV, A. N.

Use of LED disinfection units in Voronesh Providee, Veterinaria 35 no.4:71-72 Ap '58. (MIRA 11:3)

1. Machal'nik veterinarmogo otdela Voroneshskogo oblastnogo upravleniya sel'skogo khozyaystva.

(Disinfection and disinfectants)

KRYLEV, AH

#### PHASE I BOOK EXPLOITATION

SOV/4188

- Alekseyev, Valentin Petrovich, Nikolay Ivanovich Kostygov, Mikhail Georgiyevich Kruglov, Aleksey Nikolayevich Krylov, Oleg Borisovich Leonov, and Georgiy Nikolayevich Mizernyuk
- Dvigateli vnutrennego sgoraniya; opisatel'nyy kurs (Internal Combustion Engines; Descriptive Course) Moscow, Mashgiz, 1960. 451 p. 15,000 copies printed.
- Ed. (Title page): A. S. Orlin, Professor; Ed. (Inside book):
  L. I. Yegorkina; Managing Ed. for Literature on Automotive, Tractor, and Agricultural Machine Building: I. M. Bauman, Engineer; Tech. Eds.:
  B. I. Model' and T. F. Sokolova.
- PURPOSE: This textbook is intended for students at machine-building schools of higher education, and for personnel engaged in the production and operation of internal-combustion engines.
- COVERAGE: The book describes the construction and operation of all the main types of reciprocating internal-combustion engines, and of individual

Card 1/8

## Internal Combustion Engines; (Cont.)

807/4188

systems and mechanisms used in them. The book corresponds to the program of the course on "Internal-Gombustion Engines" at the Moscow Higher Technical Institute imeni N. Ye. Bauman. V. P. Alekseyev wrote chapters V and VI; N. I. Kostygov, the introduction, section 2 of chapter I, and chapters II, III and IV; M. G. Kruglov, chapter VII (except sections 40 and 42), section 57 of chapter X, and chapters XII and XIII; A.N. Krylov, chapter VIII, and sections 40 and 42 of chapter VII; O. B. Leonov, section 1 of chapter I, and chapter IX; G. N. Mizernyuk, chapters X (except section 57) and XI. The authors thank Professor D. N. Vyrubov. There are 38 references: 35 Soviet, 2 English and 1 French.

## TABLE OF CONTENTS:

For	<b>ew</b> 01	rd.				-3
Int	rodu	action				5
	I. 1. 2.	Fuel for Internal-Combustion Engines Basic information on fuel Fuel combustion in internal-combustion	engines			9 9 16
Car	1 2/	<b>'8</b>				

KRYLOV, A.N., starshiy prepodavatel'

Determining mechanical efficiency of an internal combustion engine by the running-out method. Izv.vys.ucheb.zav.; mashinostr. no.4:212-216 162. (MIRA 15:7)

1. Moskovskoye vyssheye tekhnicheskoye uchilishche imeni Baumana. (Gas and oil engines-Testing)

KRYLOV, A. P., Candidate of Biological Sciences, All-Union Academy of Agricultural Sciences imeni V. I. Lenin

"New Forms of Soya from Slowly Swollen Seed"

Doklady Vsesoyuz Ordena Lenina Skademii Sel'skhokhosyaystvennykh Nauk imeni V.I. Lenina Vol 1, 1856, pp 13-15

USSR / Human and Animal Pathology. Nervous System. Central Nervous System.

S-2

Abs Jour: Ref Zhur-Biol., No 14, 1958, 64758.

Author

Krylov, A. P. Kishinev State Medical Institute. Inst

: Morphological Changes in the Cerebrum in Primary Title

Tuberculosis.

Orig Pub: Tr. Kishinevsk. gos. med. in-ta, 1956, 5, 307-

312.

Abstract: A study was made of the cerebrum in patients who died of primary tuberculosis. The nerve cells of

the II and III layers of the cortex of the precentral area suffer the most pronounced changes. The changes in the cells of the Ammon's horn, the optic tubercle, and the hypothalamic area also have a prevailing character. However, in the hy-

Card 1/2

APPROVED FOR RELEASE 1957 147 2000 CIA-RDP86-00515R000826820012

ARYLOV, AI

24(3)

.PHASE I BOOK EXPLOITATION SOV/1643

- Avayev, Sergey Aleksandrovich, Andrey Pavlovich Krylov, and Boris Mikhaylovich Ozerskiy
- Obshchaya elektrotekhnika (General Electrical Engineering)
  Moscow, Gosenergoizdat, 1959. 447 p. 100,000 copies printed.
- Ed. (Title page): S.A. Avayev; Ed. (Inside book): M.P. Leplinskiy; Tech. Ed.: K.P. Voronin.
- PURPOSE: This book was approved by the Main Administration of Specialized Secondary Schools, Ministry of Education, USSR, as a textbook for tekhikums other than those specializing in electrical engineering.
- COVERAGE: The book contains basic information on the concept of the electric field, conductors, dielectrics, semiconductors, capacitors, d-c circuits, magnetism, single-phase a-c circuits, three-phase circuits, electrical measuring instruments, d-c machines, transformers, induction motors, converters and

Card 1/15

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

## General Electrical Engineering

## 80V/1643

amplifiers, electronic components, electric drives, electric lighting, electric power stations, substations and networks. The book also provides review questions and exercises and contains a number of laboratory experiments. The authors thank Professor B.A. Teleshev and the faculty of the Moskovskiy stankostroitel nyy tekhnikum for their assistance and Engineer M.P. Leplinskiy for editing the manuscript. There are no references.

#### TABLE OF CONTENTS:

Foreword	*
List of Units, Symbols, and Graphical Symbols	: 1
Introduction	19
Ch. I. The Electric Field  1. Electrically charged particles 2. Concept of the electric field 3. Conductors, dielectrics and semiconductors 4. Dielectrics in an electric field. Polari- sation of dielectrics. Polarisation current  Card 2/15	19 22 30 34

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

AVAYEV, Sergey Aleksandrovich, Rand. tekhn. nauk; BELOV, Vladimir
Pavlovich; ZINGMAN, Aleksandr Abramovich; MILOVIDOV,
Nikolay Nikolayevich; SIDOROV, Yuriy Pavlovich; SIMIGIN,
Petr Andreyevich; GARTUNG,S.V., retsenzent; KRYLOV, A.P.,
retsenzent; CHUGREYEVA, V.N., red.; VINOGRADOVA, G.A., tekhn.red.

[Automatisation of technological processes in the cotton industry] Avtomatizatsiia tekhnologicheskikh protsessov khlopchatobumashnoi promyshlennösti. Moskva, Gizlegprom, 1963. 279 p. (MIRA 16:11) (Cotton machinery) (Automation)

\*\*\* 1987 1 艾斯特等的特别的特殊人员特别的证明的现代的对人的分别。它是不

KRYLOV, A.P. (Kiyev); KUZ MENKO, V.A. (Kiyev); VETROV, I.Ye., inzh. (Kiyev)

larger volume of transportation with a smaller expenditure of fuel; from the experience of the Southwestern Railroad. Zhel. dor. transp. 45 no.3:70-72 Mr 163. (MIRA 16:6)

1. Nachal'nik slushby lokomotivnogo khosyaystva Tugo-Zapadnoy shelesnoy dorogi (for Krylov). 2. Nachal'nik lokomotivnogo depo Darnitsa Tugo-Zapadnoy shelesnoy dorogi (for Kus'menko).
3. Lokomotivnoye depo Darnitsa Tugo-Zapadnoy shelesnoy dorogi (for Vetrov).


(Railroads + Management) (Diesel locomotives)


BBOVMAN, M.Ya.; RIMEN, V.Kh.; BELOV, Ye.M.; KRYLOV, A.P.; VOLKOGON, G.M.

Investigation of electric power parameters in the rolling of nonferrous metals. TSvet. met. 34 no.8:60-65 Ag '61. (MIRA 14:9)

1. Yuzhno-Ural'skiy zavod tyazhelogo mashinostroyeniya (for Brovman, Rimen, Beloy. 2. Orskiy zavod obrabotki tsvetnykh metallov (for Krylov, Volkogon).

(Rolling (Metalwork)) (Nonferrous metals)





AVAYEV, Sergey Aleksanarovich; GAL'PERIN, Mikhail Moiseyevich; KKYLOV A.P., retsenzent; DIVAVIN, N.I., retsenzent; AGADZHANOVA, I.A., red.

[Fundamentals of mechanization and automation in the textile industry] Osnovy mekhanizatsii i avtomatizatsii proizvodstva v tekstil'noi promyshlennosti. Moskva, Izdvo "Legkaia industriia," 1964. 245 p. (MHRA 18:1)

T - 1 - A	Light La
	Dy, A. P., Glogovskiy, M. M., Mirchink, M. F., Nikolevskiy, N. M., and Dyy, I. A., "Scientific Bases for the Exploitation of Petroleum Deposits." Deptekhizdat, 1948, 416 pp, 3,000 copies.
T. COLLEGE	

KRYLOV, A. P.		ale vv		PA 51T92	
<b>*</b>		plaine pumps 10	"Meft Khozyay" No 1 To avoid having falling postuput of a well, it is no pressure correspondingly.  Manyo or less constant. If sealth in wells requiring	MHER/Petrol Oil W Pumps Fumps Automatic	
	ype apparatu yen under od roduced into		Khozyay"  old havin  t of a we  corre  or less o  y in well	~ # <b>6 7 4</b>	
	apparatus magr com ed into	h do	toryay" No 1 thawing falling of a well, it if it is courseponding less constant. In wells require		
		not ind	A B B L TO M	Industry	
		basic principles of new- h do not operate in deep leum Industry (Contd)	horyay" No 1  having falling plastic pressure affect the of a well, it is necessary to decrease the factor correspondingly. Thus the depression remain less constant. This is brought about fairly in wells requiring deep pumping. Author ex-	leem Industry 11s Operating Couplings,	
	•	ete ble	14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15	up l du	
		(Contd)	plastic pressure affect necessary to decrease to the depression. This is brought about in this is brought about in	89	
	rareign:	new-type leep well	pressure the dep trought	<b>&gt;</b>	
	nat.	walls	depr depr	₽l	÷ .
	, i 🔑 🚅	cou	affect mease t ession about f	Ja Erylor,	
J	T was	coupling for 51792	affect the ease the fi ession remain bout fairl		
25 T.	Ä.	ling for his new- 51792 Jan 1948	the face fairly	1948 3. V.	
	1.0	7	EV		
erson secondence secondence		MAIN BONNEY TO	LI BOARS DIN E WATER		Terra di Sa

MURAVEV, I. M. and KRYLOV, A. F.

Eksploatatsiia Heftianykh Mestorozhdenii (Oil Field Exploitation), 775 p., Moscow and Leningrad, 1949.

APPROVED FOR RELEASE: 06/14/2000 CIA-RDP86-00513R000826820012-2"

